


SEARS & ZEMANSKY'S  
**College  
Physics**  
*9th edition*

PowerPoint Lectures

**1** **Models, Measurements  
and Vectors**

Lectures by **James L. Pazun**

*Young*



## *Goals for Chapter 1*

- To know standards and units and be able to do unit conversions.
- To express measurements and calculated information with the correct number of significant figures.
- To be able to add vectors.
- To be able to break down vectors into  $x$  and  $y$  components.

# Measurement

- Physics is an experimental science.
  - Observe phenomena in nature.
  - Make predictions.
    - Models
    - Hypothesis
    - Theories
    - Laws

# *Units of measurement*

- Cultural
  - “cubit”, “span”, “foot”, “mile”
  - Changes with time and location
- 1889 by the General Conference on Weights and Measures
  - Systéme International

# *There are three fundamental S.I. units.*

- Time

- Second (s)

- Length

- Meter (m)

- Mass


- Kilogram (kg)

# *The Second*

- Originally tied to the length of a day.
- Now, exceptionally accurate.
  - Atomic clock
  - 9,192,631,770 oscillations of a low-energy transition in Cs
  - In the microwave region

## The meter - Figure 1.3a

The meter was originally defined as 1/10,000,000 of this distance.




## *The meter – more recently*

- Now tied to Kr discharge and counting a certain number of wavelengths.
- Exceptionally accurate, in fact redefining  $c$ , speed of light.
- New definition is the distance that light can travel in a vacuum in  $1/299,792,458$  s.
- So accurate that it loses only 1 second in 30 million years.



*Figure 1.3b*



# *The kilogram*

- The reference cylinder is kept in Sevres, France.
- A more modern, atomic reference is hoped for, waiting for a precise atomic technique.

# *The reference kilogram - Figure 1.3c*



## *You can adjust the fundamental units.*

- Imagine trying to measure the distance from San Francisco, CA to Charlotte, NC in meters.
- Instead, we attach prefixes to the units that adjust their size by powers of ten.
  - In this case kilo (or “ $\times 10^3$ ”) would likely be chosen.
  - The distance would then be reported as 4,621km or 4.6Mm.
    - Getting good at converting units will be very important to us.
    - Use Table 1.1.

# Table 1.1

**TABLE 1.1 Prefixes for powers of ten**


Power of ten	Prefix	Abbreviation
$10^{-18}$	atto-	a
$10^{-15}$	femto-	f
$10^{-12}$	pico-	p
$10^{-9}$	nano-	n
$10^{-6}$	micro-	$\mu$
$10^{-3}$	milli-	m
$10^{-2}$	centi-	c
$10^3$	kilo-	k
$10^6$	mega-	M
$10^9$	giga-	G
$10^{12}$	tera-	T
$10^{15}$	peta-	P
$10^{18}$	exa-	E

# The powers of ten are dramatic – Figure 1.4

$10^{21}$  m  
A galaxy




$10^9$  m  
Our sun




$10^7$  m  
The earth




$10^0$  m  
You



$10^{-5}$  m  
Red blood cells



$10^{-9}$  m  
Individual atoms



$10^{-14}$  m  
An atomic nucleus

# Conversions

- “Practice, practice, practice.”
  - Problems will come in many different styles of measurement but you will ultimately need to get back to (m), (kg), and (s) if your answers are going to fit other calculations.
  - You’ll need to overcome two hurdles:
    - Derived units
    - English → S.I.
  - Let’s examine each.

## *Derived units*

- The science will wait until we find it in a subsequent chapter. First, we should examine some units from those chapters.
  - Imagine you need to work with energy.
  - The unit for energy is Joule (J) and it's built from other units: in this case  $\text{kg m}^2/\text{s}^2$ .
  - Having mass in g or distance in cm ... errors like that will destroy your answer.




- In the United States, we often encounter measurements in miles, feet, pounds, quarts and gallons.
- It's useful to memorize one conversion for displacement, one for volume, and one for mass. Everything else could be reached by converting orders of magnitude.
- Mass is tricky. English units assume standard earth gravity and relate kg to lb even though kg is a mass and lb is a force.

- Personal favorites for English to SI are listed below.
  - Displacement: 2.54 centimeters = 1 inch.
  - Mass: 454 grams = 1 pound.
  - Volume: 1 liter = 1.06 quarts.
- You could certainly choose a different set.

# Precision and Significant Figures

- Your measurement tools have limitations and your reported results need to reflect those limitations.



## *“Am I significant?”*

- At the risk of making data feel unwanted, we dare not report everything our calculator tells us. (Please forgive the anthropomorphism.)
  - Try this. Divide 10 (one SF) by 3 (one SF) and your calculator will tell you 3.333333333.
  - If you report this answer, the reader will believe you have measured carefully to billionths of the unit you are using.
  - What can happen? It’s possible that bolt holes will fail to line up.

*This could happen.*




# Vector addition


- In the “world of vectors”  $1+1$  does not necessarily equal 2.

- Graphically?

Vectors  $\vec{A}$  and  $\vec{A}'$  are equal because they have the same length and direction.




This vector is different from  $\vec{A}$ ; it points in the opposite direction.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings

To find the sum of these three vectors ...




(a)

we could add  $\vec{A}$  and  $\vec{B}$  to get  $\vec{D}$  and then add  $\vec{C}$  to  $\vec{D}$  to get the final sum (resultant)  $\vec{R}$ , ...




(b)

or we could add  $\vec{B}$  and  $\vec{C}$  to get  $\vec{E}$  and then add  $\vec{A}$  to  $\vec{E}$  to get  $\vec{R}$ , ...




(c)

or we could add  $\vec{A}$ ,  $\vec{B}$ , and  $\vec{C}$  to get  $\vec{R}$  directly, ...




(d)

or we could add  $\vec{A}$ ,  $\vec{B}$ , and  $\vec{C}$  in any other order and still get  $\vec{R}$ .




(e)

*Or, decompose the vectors into components then solve.*



(a)




(b)

# Just in case your trigonometry is rusty, let's review.

Try example 1.6 to check yourself.

**First draw your axes. Make them big enough.**



**Then draw in the displacement vectors and their components.**

**Be sure to put an arrow over each vector symbol, and add x and y subscripts to the component symbols.**