Instructions:

Part II- do on page 4 of scantron mini blue book

Multiple Choice: Choose the single best response to each question.

- 1. Tides in the earth's oceans are caused by
  - a. only the earth's gravity d. both sun and moon's gravity
  - b. only the moon's gravity e. both earth and moon's gravity
  - c. only sun's gravity
- 2. If you double the mass of the earth, the gravitational force of the sun on the earth will be changed to \_\_\_\_\_\_ times its original value
  a. 0.25 b. 0.50 c. 1 d. 2 e. 4

3. A planet is in orbit around a star. The value for g of the star at the orbit of the planet is 5 m/s<sup>2</sup>. The star suddenly collapses to a new diameter that is one half of its original diameter. The mass of the star does not change during the collapse. The value of g at the planet's orbit after collapse is \_\_\_\_\_m/s<sup>2</sup>.

a. 1.25
b. 25.0
c. 5.00
d. 10.0
e. 20.0

You have arrived at a planet that has a radius the same as the earth's radius and a mass that is three times the mass of the earth. Use this information to answer numbers 4 through 6.

- 4,5. Find the value for g, the acceleration due to gravity on the planet's surface, for this planet in terms of g on earth. In other words  $g_p = xg_E$ . Find x. a. 1/9 b. 1/3 c. 1 d. 3 e. 9
  - 6. Find the value for g, the acceleration due to gravity on the planet's surface in m/s<sup>2</sup> a. 1.09 b. 3.27 c. 9.80 d. 29.4 e. 88.2

The next planet on your trip has a radius 2 times the Earth's radius and a mass the same as the mass of earth. Use this information to answer numbers 7 through 9.

- 7,8. Find the value for g, the acceleration due to gravity on the planets' surface, for this<br/>planet in terms of g on earth. In other words  $g_p = xg_E$ . Find x.<br/>a.  $\frac{1}{4}$ b.  $\frac{1}{2}$ c. 1d. 2e. 4
  - 9. Find the weight in newtons of a 5 kg object on the planet's surface. a. 12.3 b. 24.5 c. 49.0 d. 98.0 e. 196
  - 10. People in orbit around the earth in the space shuttle appear to be weightless because they
    - a. are above the air d. are in free fall
    - b. are far from earth e. don't feel the gravitational force in orbit
    - c. are moving rapidly

11. Bullet 1 is dropped (by hand) at the same time that bullet 2 is fired horizontally from a rifle. Which bullet lands first? Neglect air resistance.

a. 1 b. 2 c. neither, they land at the same time

12. A baseball is thrown horizontally with a speed of 15 m/s. Once the ball has left the thrower's hand, the horizontal acceleration of the ball is \_\_\_\_\_m/s<sup>2</sup>. neglect air resistance

a. 0 b. 5.2 c. 9.8 d.15.0 e. 24.8

13. A baseball is thrown horizontally with a speed of 15 m/s. Once the ball has left the thrower's hand, the vertical acceleration of the ball is  $_{m/s^2}$ . neglect air resistance

a. 0 b. 5.2 c. 9.8 d. 15.0 e. 24.8

A 0.2 kg arrow is shot horizontally with a speed of v from a cliff 15 meters above the level ground below the cliff. The arrow lands 75 m from the base <u>of the cliff as shown</u> v to the right. Use this information to answer numbers 14 through 17.

- 14,15. How many seconds is the arrow in the air?
   15 m

   a. 1.75
   b. 2.17
   c. 3.06
   d. 3.91
   e. 15.3
- 16,17. Find the speed v in m/s of the arrow when it leaves the bow. ↓ a. 4.90 b. 19.2 c. 24.5 d. 34.6 e. 42.9

A car is following a curved path from point 1 to point 3 as shown to the right. Use this information to answer numbers 18 through 23.

- 18. The car's speed is increasing at point 1 Which arrow shows the direction of the car's velocity vector?
- 19. The car's speed is constant at point 2. Which arrow shows the direction of the car's velocity vector?
- 20. The car's speed is decreasing at point 3. Which arrow shows the direction of the car's velocity vector?
- 21. The car's speed is increasing at point 1. Which arrow shows the direction of the car's acceleration vector?
- 22. The car's speed is constant at point 2. Which arrow shows the direction of the car's acceleration vector?
- 23. The car's speed is decreasing at point 3. Which arrow shows the direction of the car's acceleration vector?



75 m

Name:	Name:					Physics5-Test2					Date	
24. 7	The ce a.	entripet direct	al force ion of tra	causes a vel	an objec b. spec	et's ed of tra	to vel	change c. noth	ing			
25,2	25,26. If your car goes around a turn with a speed $\mathbf{v}_1$ , it takes a centripetal force $F_1$ keep the car going around the turn. What force $F_2$ is required if you go around the curve with a speed $\mathbf{v}_2$ that is 1.5 times $\mathbf{v}_1$ ?											
	a.	$F_1$		b. 1.50	) F <sub>1</sub>	c. 2.25	$F_1$	d. 3.00	$F_1$	e. 4.00 F	1	
A 0.1 rock thou	A 0.3 kg rock is being whirled at the end of a string in a circle of radius 2.0 m. the rock makes 4 revolutions every second. Use this information to answer numbers 27 though 30.											
27,2	8. Fin a. 1	d the sj 2.00	peed of t	he rock b. 3.14	t in m/s. 1	c. 6.28		d.12.6		e. 50.3		
29. H	ind tl a.	ne centr 2.00	ripetal ac b. 4.93	ccelerat	tion of t c. 19.7	he rock	in m/s <sup>2</sup> . d. 79.4		e. 1,26	0		
30. H	ind tl a.	ne centr 0.600	ripetal fo	orce in 1 b. 1.48	N on the 3	e rock c. 5.91		d. 23.8		e. 378		
31.7	<ul><li>31. The net force on a skydiver is greatest when she</li><li>a. first steps out of the airplane</li><li>b. has reached terminal velocity</li><li>c. is halfway between a. and b.</li></ul>											
32. 1	<ul><li>32. The acceleration of a skydiver is least when she</li><li>a. first steps out of the airplane</li><li>b. has reached terminal velocity</li><li>c. is halfway between a. and b.</li></ul>											
Use the following information to work problems 33 though 38. A man is measuring the acceleration of an express elevator in the Empire State Building by reading his old bathroom scale that he is standing on while riding the elevator. Before the elevator starts moving, the scale reads 185 lb.												
33,34,35. While starting to move, the scale reads 165 lb. find the magnitude acceleration of the elevator in $ft/s^2$ .												
	a.	0.289	b. 3.46		c. 5.78		d. 28.5		e. 32.0			
36.	What a.	is the c up.	lirection b. dow	of the an	accelera c. ther	tion? e is no v	vay to to	ell				
37,38. Later the elevator is traveling at a constant speed of 15 ft/s. Find the reading on												
	the a.	scale 1 165	b. 170	S.	c. 185		d. 200		e. 205			

- 39,40,41. How many feet in altitude would a skydiver have to be in order to be able to fall through the air for 25 seconds? Neglect air resistance.
  - a. 123 b. 400 c. 800 d. 37,062 e. 10,000
- 42. Which of the following statements is correct?
  - a. 1 kg isn't related to 2.2 lb d. 1 kg equals 2.2 lb
  - b. 1kg has a mass of 2.2 lb e. none of these
  - c. 1 kg weighs 2.2 lb

Identify the following as either: free fall (mark a) of not free fall (mark b)

- 43. A feather dropped from two meters above the earth
- 44. A rock dropped from two meters above the earth
- 45. A satellite in orbit around the earth
- 46. The space shuttle during re-entry through the earth's atmosphere

Part 2

- An object weighs 15 N when on the surface of the earth. It is then moved to a distance 3.5 times the earth's radius, (measured from the center of the earth). Find: the weight of the object at this distance. Ans: 1.22 N
- The mass of Venus is 4.90x10<sup>24</sup> kg and its distance from the sun is 1.08x10<sup>11</sup> m. It takes 1.94x10<sup>7</sup> s to complete an orbit. Find: the gravitational force exerted on Venus by the Sun. Ans: 5.54x10<sup>22</sup> N
- 3. Reproduce the following drawing on page 4 of your scantron mini blue book and indicate the vectors for the following at points 1 and 2 on the projectile path provided.
  - a. acceleration b. vertical velocity component
  - c. horizontal velocity component

